

In the name of God

Antepartum Fetal Assessment

Dr. Mahboobe Sadat Miri

Faculty member of Kamali Hospital

Summer 2024

Types of fetal assessment

☐ fetal movement counts (“kick counts”)


Fetal movement counting (also called “kick counts”) is a test that the woman can do at home. There are different ways kick counts can be done. the health care professional will tell her how often to do it and when to notify him or her.

Daily fetal movement counting, such as the Cardiff “count-to-ten” method using kick charts, is a way of screening for fetal well-being, by which a woman counts daily fetal movements to assess the condition of her baby.

INSTRUCTIONS

- ❖ Choose one period during the day to count. should choose a time when the baby is normally active.
- ❖ Count at the same time every day.
- ❖ chart how long it takes to reach 10 movements.
- ❖ Count all recognizable movements. This may be a kick, a punch, rolling, stretching, etc. If you feel a short flurry of kicks, count that as one movement. Do not count hiccups.

A way to assess intrauterine well-being in which the expectant woman records fetal movement during her usual activities. There should be at least 10 movements within a 2-hour period; if fewer than 10 movements are perceived, further medical evaluation is needed.



FETAL KICK COUNT

1	Start Time					
	Stop Time					
2	Start Time					
	Stop Time					
3	Start Time					
	Stop Time					
4	Start Time					
	Stop Time					
5	Start Time					
	Stop Time					
6	Start Time					
	Stop Time					
7	Start Time					
	Stop Time					
8	Start Time					
	Stop Time					
9	Start Time					
	Stop Time					
10	Start Time					
	Stop Time					

Mark an x in the coloured box every time you feel you feel a movement. You can stop when you reach 10 movements. Opinions vary on how long it should take you to feel 10 movements. Talk to your Doctor.

Non Stress Test (NST)

- ▶ Looks to accelerations in FHR as an indication of fetal well-being; a fetus that is NOT acidotic or neurologically depressed will have a HR that temporarily accelerates with fetal movement

REACTIVE: ≥ 2 accelerations within 20 minutes

NONREACTIVE: < 2 accelerations in 40 minutes

Accelerations ≥ 32 weeks: 15 BPM above baseline + lasting $\geq 15s^{4-5}$

Accelerations < 32 weeks: 10 BPM above baseline + lasting $\geq 10s$

If it is believed that there is an absence of accelerations 2/2 fetal sleep cycles, **vibroacoustic stimulation**-elicited accelerations are a valid prediction of FWB⁶⁻⁹

- ▶ Advantages: doesn't require an IV, oxytocin, or contractions
 - ▶ no contraindications

Non Stress Test: Significance

▶ Non reactivity is associated with:

▶ Fetal distress in labor (5x)

▶ Low 5 min. Apgar scores (6x)

Increased Fetal Death Rate (7-12x)

Decelerations during a Reactive NST:

Increased distress in labor (2 – 3X)

IUGR (8 – 12X)

▶ Fetal Death (5X)

The non-stress test

The non-stress test measures the fetal heart rate in response to fetal movement over time. The term “non-stress” means that during the test, nothing is done to place stress on the fetus.

This test may be done in the health care professional’s office or in a hospital. The test is done while woman are reclining or lying down and usually takes at least 20 minutes. A belt with a sensor that measures the fetal heart rate is placed around the abdomen. The fetal heart rate is recorded by a machine.

NON-STRESS TEST (NST)





Results mean :

If two or more accelerations occur within a 20-minute period, the result is considered reactive or “reassuring.” A reactive result means that for now, it does not appear that there are any problems. A nonreactive result is one in which not enough accelerations are detected in a 40-minute period. It can mean several things. It may mean that the fetus was asleep during the test. If this happens, the test may last 40 more minutes, or the fetus may be stimulated to move with sound projected over the mother’s abdomen. A nonreactive result can occur if the woman has taken certain medications. It also can mean that the fetus is not getting enough oxygen.

Biophysical profile

A biophysical profile (BPP) may be done when results of other tests are non-reassuring. It uses a scoring system to evaluate fetal well-being in these five areas:

1. Fetal heart rate
2. Fetal breathing movements
3. Fetal body movements
4. Fetal muscle tone
5. Amount of amniotic fluid

TABLE 20-2. Components and Scores for the Biophysical Profile

Component	Score 2	Score 0
Nonstress test ^a	≥2 accelerations within 20–40 min	0 or 1 acceleration within 20–40 min
Fetal breathing	≥1 episode of rhythmic breathing lasting ≥30 sec	<30 sec of breathing
Fetal movement	≥3 discrete body or limb movements	<3 discrete movements
Fetal tone	≥1 episode of extremity extension and subsequent return to flexion	0 extension/flexion events
Amnionic fluid volume ^b	A pocket of amnionic fluid that measures at least 2 cm in two planes perpendicular to each other (2 × 2 cm pocket)	Deepest single vertical pocket ≤2 cm

^aMay be omitted if all four sonographic components are normal.

^bFurther evaluation warranted, regardless of biophysical composite score, if deepest vertical amnionic fluid pocket ≤2 cm.

TABLE 20-3. Interpretation of Biophysical Profile Score

Biophysical Profile Score	Interpretation	Recommended Management
10	Normal, nonasphyxiated fetus	No fetal indication for intervention; repeat test weekly
8/10 (Normal AFV)	Normal, nonasphyxiated fetus	No fetal indication for intervention; repeat test weekly
8/8 (NST not done)		
8/10 (Decreased AFV)	Chronic fetal asphyxia suspected	If ≥ 36 weeks, deliver If < 36 weeks, monitor per institution's protocol
6 (Normal AFV)	Equivocal	If ≥ 37 weeks, deliver If < 37 weeks and normal fluid, repeat test in 24 hours If repeat test > 6 , monitor per institution's protocol
6 (Decreased AFV)	Possible fetal asphyxia	If 36–37 weeks, deliver If < 36 weeks, monitor per institution's protocol
4	Probable fetal asphyxia	If ≥ 32 weeks, deliver If < 32 weeks, individualize management based on maternal and fetal conditions
0 or 2	Almost certain fetal asphyxia	Deliver

AFV = amnionic fluid volume; NST = nonstress test.

Adapted from American College of Obstetricians and Gynecologists, 2021a; Liston, 2018; Manning, 2018.

Activate Windows
Go to Settings to activate Windows.

Contraction stress test

A **contraction stress test** (CST) is performed near the end of pregnancy (34 weeks' gestation) to determine how well the fetus will cope with the contractions of childbirth. The aim is to induce contractions and monitor the fetus to check for heart rate abnormalities using a cardiotocograph. A CST is one type of antenatal fetal surveillance technique.

TABLE 20-1. Criteria for Interpretation of the Contraction Stress Test

Negative: no late or significant variable decelerations





Positive: late decelerations following 50% or more of contractions (even if the contraction frequency is fewer than three in 10 minutes)

Equivocal-suspicious: intermittent late decelerations or significant variable decelerations

Equivocal-hyperstimulatory: fetal heart rate decelerations that occur in the presence of contractions more frequent than every 2 minutes or lasting longer than 90 seconds

Unsatisfactory: fewer than three contractions in 10 minutes or an uninterpretable tracing

Contraction Stress Test





Uses

The CST is used for its high negative predictive value. A negative result is highly predictive of fetal wellbeing and tolerance of labor. The test has a poor positive predictive value with false-positive results in as many as 30% of cases. A positive CST indicates high risk of fetal death due to hypoxia and is a contraindication to labor. Patient's obstetricians usually consider operative delivery in such situations.

Contraindication

This "stress test" is usually not performed if there are any signs of

1. Premature birth
2. placenta previa, vasaprevia
3. Cervical incompetence
4. Multiple gestation.
5. previous classic cesarean section
6. Other contraindications include but are not limited to previous uterine incision with scarring, previous myomectomy entering the uterine cavity, and PROM. Any contraindication to labor is contraindication to CST.

CST is performed weekly as the fetus is assumed to be healthy after a negative test and should remain so for another week. This test is done in hospital or clinic setting. External fetal monitors are put in place and then either nipple stimulation or IV Pitocin (oxytocin) is used to stimulate uterine contractions.

Nipple stimulation

This is a procedure that relies on endogenous release of oxytocin following nipple stimulation, and is conducted by the patient.


The nurse instructs the patient on the procedure, as follows. One nipple is massaged gently through clothing until a contraction begins, or for a maximum of 2 minutes. If at least 3 contractions in 10 minutes is not achieved, then the patient rests for 5 minutes and the other nipple is stimulated.

Oxytocin challenge test (OCT)

If adequate contractions (at least 3 in 10 minutes) cannot be achieved with nipple stimulation, an oxytocin challenge test may be performed.

It involves the intravenous administration of exogenous oxytocin to the pregnant woman. The target is to achieve around three contractions every ten minutes

oxytocin challenge test



OCT (oxytocin challenge test)

OCT positive

The contraction stress test helps the health care professional see how the fetal heart rate reacts when the uterus contracts. The contraction stress test sometimes is used if other test results are positive or unclear.

In this test, belts with sensors that detect the fetal heart rate and uterine contractions are placed across the abdomen. To make the uterus contract mildly, pregnant woman may be asked to rub her nipples through clothing or may be during given oxytocin.



Umbilical Artery


- Doppler is very important in the management of IUGR
- Power Spectral Density (PSD) as a routine use of detecting IUGR is not proven to be better than fetal biometric parameters.
- It has, however, been proven to be a good indicator of fetal outcome
- Umbilical artery (UA) Doppler in suspected IUGR is performed.
- If UA is normal, then no other Doppler investigations need to be performed.
- If UA is abnormal, then other vessels should be evaluated.

- With placental insufficiency, placental impedance is higher than normal and results in diastolic abnormalities such as decreased, absent, or reversed end-diastolic flow in fetal vessels.
- These diastolic abnormalities in flow are best characterized by the standard Doppler indices (S/D ratio, RI, and PI).


- The diastolic flow reversal in umbilical arteries signifies severe placental insufficiency and increased placental vascular resistance, which is bad news for the fetus.

- Normally progressive decrease in resistance during the course of pregnancy.
- $RI < 0.8$
- Systolic/diastolic ratio > 3.0 is abnormal after 30 weeks
- Ratios are higher if measured closer to fetal cord insertion, so measurements should be taken in UA close to placental cord insertion site
- Absent or reversed diastolic flow in umbilical artery is ALWAYS an ominous sign


Abnormal UA Doppler waveform depends on the severity of disease.




Normal pregnancy



Reduced end diastolic velocity




Absent end diastolic velocity



Reversed end diastolic velocity

Normal MCA in the fetus




- IUGR fetuses may be experiencing chronic hypoxemia (decreased blood oxygen concentration) .
- With fetal hypoxemia, there is increased blood flow to vital organs (brain and myocardium) and reduced flow to the gastrointestinal tract and kidneys.
- Thus, the PSV of a hypoxic fetus will show an increase in the MCA velocity.

MCA and ANEMIA

Middle Cerebral artery (MCA) Doppler can help determine likelihood of fetal anemia.

MCA is examined close to its origin from the internal carotid artery.

An abnormal MCA.



Uterine Artery

- Low resistance in second and third trimesters is normal
- Appearance of a 'notch' in Doppler waveform is abnormal and thought to predict insufficiency & IUGR

- IVC Doppler waveforms can be obtained from a coronal plane of the chest and abdomen. In this view, the IVC can be imaged as it enters into the right atrium, joined by the ductus venosus and the left hepatic vein. The IVC waveforms are triphasic in shape, with the first phase corresponding to ventricular systole, the second phase to early diastole, and the third phase to late diastole or the atrial kick.

Indications for Antepartum Testing

Maternal

Chronic hypertension
Pregestational DM
SLE
Antiphospholipid syndrome
Hemoglobinopathies
Cyanotic heart disease
Cardiomyopathy
Cystic fibrosis
Restrictive lung disease
Chronic renal disease
Hyperthyroidism
In vitro fertilization
Substance abuse
Chemotherapy (current)
Prepregnancy BMI ≥ 35
Maternal age >35

Pregnancy-related


Gestational hypertension
Preeclampsia
Insulin-requiring gestational DM
Oligohydramnios
Polyhydramnios
Postterm pregnancy
Prior stillbirth
Isoimmunization
Cholestasis
Velamentous cord insertion
Single umbilical artery

Fetal

Fetal-growth restriction
Decreased fetal movement
Multifetal gestation

BMI = body mass index; DM = diabetes mellitus;
SLE = systemic lupus erythematosus.

Activate Windows
Go to Settings to activate Windows.

A top-down photograph of a white card with the words "Thank you" written in a vibrant purple, cursive script. The card is placed on a light-colored, marble-textured surface. To the left of the card is a small bouquet of purple bell-shaped flowers with green leaves. To the right of the card lies a black pen with a white band decorated with black stars. A string of red and white braided beads is coiled around the pen and extends towards the top right corner. In the bottom right corner, a portion of a gift wrapped in white paper with grey polka dots and tied with the same red and white braided string is visible.

Thank
you